NASA/TM-2000-209891, Vol. 157



Technical Report Series on the Boreal Ecosystem-Atmosphere Study (BOREAS)

Forrest G. Hall and Shelaine Curd, Editors

Volume 157 BOREAS TE-9 NSA Photosynthetic Response Data

Qinglai Dang, Lakehead University, Thunder Bay, Canada Hank Margolis and Marie Coyea, Université Laval, Sainte-Foy, Quebec, Canada

National Aeronautics and Space Administration

Goddard Space Flight Center Greenbelt, Maryland 20771

	····	
		·
	A 11 - 1. 1 . C	
	Available from:	
NASA Center for AeroSpace Information 7121 Standard Drive Hanover, MD 21076-1320 Price Code: A17		National Technical Information Service 5285 Port Royal Road Springfield, VA 22161 Price Code: A10

BOREAS TE-9 NSA Photosynthetic Response Data

Qinglai Dang, Hank Margolis, Marie Coyea

Summary

The BOREAS TE-9 team collected several data sets related to chemical and photosynthetic properties of leaves. This data set describes (1) the response of leaf and shoot-level photosynthesis to ambient and intercellular CO₂ concentration, temperature, and incident PAR for black spruce, jack pine, and aspen during the three IFCs in 1994 in the NSA; (2) the response of stomatal conductance to vapor pressure difference throughout the growing season of 1994; and (3) a range of shoot water potentials (controlled in the laboratory) for black spruce and jack pine. The data are available in tabular ASCII files

Table of Contents

- 1) Data Set Overview
- 2) Investigator(s)
- 3) Theory of Measurements
- 4) Equipment
- 5) Data Acquisition Methods
- 6) Observations
- 7) Data Description
- 8) Data Organization
- 9) Data Manipulations
- 10) Errors
- 11) Notes
- 12) Application of the Data Set
- 13) Future Modifications and Plans
- 14) Software
- 15) Data Access
- 16) Output Products and Availability
- 17) References
- 18) Glossary of Terms
- 19) List of Acronyms
- 20) Document Information

1. Data Set Overview

1.1 Data Set Identification

BOREAS TE-09 NSA Photosynthetic Response Data

1.2 Data Set Introduction

The response of photosynthesis to ambient CO₂ concentration, temperature, light (Photosynthetically Active Radiation (PAR)), vapor pressure difference (VPD), and shoot water potential was investigated as part of an effort to construct the response surfaces of photosynthesis to different environmental factors. Samples were taken from three forest types: jack pine (Pinus banksiana Lamb.), black spruce (Picea mariana Mill. B.S.J.P), and aspen (Populus tremuloides Michx.) in the BOReal Ecosystem-Atmosphere Study (BOREAS) Northern Study Area (NSA) during each of the three Intensive Field Campaigns (IFCs) in 1994. Measurements were taken under controlled environmental conditions in the laboratory using an open gas exchange system in differential mode. Photosynthesis and related parameters all are expressed on a hemisurface area basis. The shape factors for leaf area calculation are 4 and 4.59, respectively, for black spruce and jack pine.

1.3 Objective/Purpose

This data set was collected and prepared to provide the response curves of photosynthesis to (1) ambient and intercellular CO₂ concentration, (2) temperature, and (3) PAR in jack pine, black spruce, and aspen in the NSA using a cut-branch technique. Additional data sets were collected and prepared to provide the response curves of photosynthesis and stomatal conductance to water vapor pressure difference for jack pine and black spruce to provide the response of photosynthesis to shoot water potential in jack pine and black spruce in the NSA.

1.4 Summary of Parameters

Net photosynthesis, ambient and intercellular CO₂ concentration, transpiration, stomatal conductance, temperature, PAR, VPD, water potential.

1.5 Discussion

The response of photosynthesis to ambient CO₂ concentration, temperature, and light (PAR), the response of stomatal conductance to VPD; and the response of photosynthesis and stomatal conductance to shoot water potential were investigated as part of an effort to construct the response surfaces of photosynthesis to different environmental factors. Samples were taken in the NSA during each of the three IFCs in 1994 from three forest types: old jack pine (OJP), old black spruce (OBS), and old aspen (OA). Measurements were taken under controlled environmental conditions in the laboratory using an open gas exchange system in differential mode. Photosynthesis and related parameters all are expressed on a hemisurface area basis. The shape factors for leaf area calculation are 4 and 4.59, respectively, for black spruce and jack pine.

1.6 Related Data Sets

BOREAS TE-09 NSA Photosynthetic Capacity and Foliage Nitrogen Data BOREAS TE-09 PAR and Leaf Nitrogen Data for NSA Species BOREAS TE-09 NSA in situ Diurnal Gas Exchange of Boreal Forest Species

2. Investigator(s)

2.1 Investigator(s) Name and Title

Dr. Hank Margolis, Associate Professor

2.2 Title of Investigation

Relationship Between Measures of Absorbed and Reflected Radiation and the Photosynthetic Capacity of Boreal Forest Canopies and Understories

2.3 Contact Information

Contact 1:

Dr. Hank Margolis Universite Laval Faculte de foresterie et de geomatique Pavillon Abitibi-Price Sainte-Foy, Quebec Canada G1K 7P4 (418) 656-7120 Hank.margolis @sbf.ulaval.ca

Contact 2:

Dr. Qinglai Dang Lakehead University Faculty of Forestry 955 Oliver Road, Thunder Bay Canada P7B 5E1 (807) 343-8507 (807) 343-8116 (fax) Qinglai.Dang@flash.lakeheadu.ca

Contact 3:

Shelaine Curd Raytheon ITSS Code 923 NASA GSFC Greenbelt, MD 20771 (301) 286-2447 (301) 286-0239 (fax) shelaine.curd@gsfc.nasa.gov

3. Theory of Measurements

During the process of photosynthesis, CO₂ is assimilated by green leaves (photosynthesis) while H₂O is released into the atmosphere (transpiration). The amount of water released and the amount of CO₂ absorbed can be determined by comparing the concentrations of water vapor and CO2 in the air moving into the leaf cuvette and those in the air moving out of the cuvette at a known flow rate. The concentrations of CO2 and water vapor in both incoming and outgoing air streams can be measured using an infrared gas analyzer (IRGA). The rates of net photosynthesis and transpiration are calculated from the difference in the concentrations of CO₂ and water vapor between the input and the output from the leaf cuvette. Stomatal conductance is calculated from transpiration rate and the water vapor gradient between the intercellular space and the bulk air in the cuvette.

The water in the xylem is under tension. When the stem of the branch is cut, the water will retreat from the cut surface. When the cut branch is enclosed in the pressure chamber with the cut surface extruding and pressurized gradually, the xylem water will come back to the cut surface when the pressure is equal to the water potential of the shoot. The pressure inside the pressure chamber, and

thus the water potential of the shoot, can be read from a pressure gauge.

4. Equipment

4.1 Sensor/Instrument Description

LI-COR 6262 IRGA, thermocouples, balance, Decagon AgVision root and leaf analysis system, PMS Model 610 pressure chamber.

4.1.1 Collection Environment

Values of major environmental variables are given in the data set for each individual measurement. Upper-canopy branch samples were harvested using a shotgun and were immediately recut under water. The samples were then transported to the laboratory for gas exchange measurement. The cut surfaces of the branches were submerged in water during transport (30 to 50 min) and in the laboratory. Measurements for each species generally took 6 to 10 hours.

All samples were kept in the dark, but prior to measurement, samples were exposed to saturated light for 2 hours to induce stomatal opening and photosynthetic activity. To test photosynthetic

response to CO₂ concentration changes, steady-state readings were taken at each CO₂ level. An independent set of two samples was used for each two CO₂ levels. Ambient CO₂ varied from 50 to 900 ppm.

To keep a continued supply of water to the branch, the cut surface was kept in contact with water during the entire course of measurement. Saturated light was supplied using two 1,000-watt high-pressure sodium lamps. When ambient CO_2 was varied, other environmental conditions were as follows: temperature = 20 +/- 0.5 °C; VPD = 0.7 +/- 0.1 kPa; CO_2 = 360 +/- 20 ppm.

To test photosynthetic response of varied temperature, the air temperature inside the leaf cuvette was controlled using a radiator that was driven by a temperature-controlled water bath. The vapor pressure inside the cuvette was controlled by passing water vapor-saturated air through a condenser whose temperature was controlled using another water bath.

An independent set of two branches was used for each temperature. The branches were from four different trees and were mixed randomly.

Saturated light for the measurement was supplied using two 1,000-watt high- pressure sodium lamps. VPD of the air was controlled at a relatively constant level except at temperatures below $10\,^{\circ}$ C, when there were some technical difficulties in getting a low enough vapor pressure to maintain the desired VPD. Input CO₂ concentration was controlled at $360 \, (+/-15 \, \text{ppm})$.

To test photosynthetic response of varied PAR, the measurements started from the highest PAR level and proceeded to darkness. Steady-state readings were taken at each light level. The light source was two 1,000-watt high-pressure sodium lamps. Different levels of light were achieved by using different neutral density filters. The environmental conditions inside the leaf cuvette were as follows: temperature 20 +/- 0.5 °C; VPD 0.7 +/- 0.2 kPa; CO₂ 360 +/- 15 ppm.

To test stomatal conductance of varied VPD, all samples were kept in the dark. The samples to be measured, however, were exposed to saturated light for 2 hours prior to measurement to induce stomatal opening and photosynthetic activity.

The stability and reliability of the cut-branch technique were tested. Stable measurements for at least 24 hours are possible. Steady-state readings were taken at each VPD level.

An independent set of two samples was used for each VPD level. To keep a continued supply of water to the branch, the cut surface was kept in contact with water during the entire course of measurement.

Different VPD levels were achieved by regulating the water vapor pressure of the input air stream to the leaf cuvette. Saturated light was supplied using two 1,000-watt high-pressure sodium lamps. The CO_2 concentration in the input air was 360 (+/- 15 ppm). Measurements were taken at three temperatures (15, 25, and 35 °C) in IFC-1, two temperatures (25 and 35 °C) in IFC-2, and at 25 °C only in IFC-3.

In the laboratory, the branches were taken out of the water, the cut surfaces of the branches were dried and sealed using silicon grease. The branches were then exposed to light and let transpire freely. At certain time intervals, the gas exchange of the branches (two at a time) was measured. The water potential of the branches was measured immediately after the gas exchange measurement.

Gas exchange was measured at saturated light conditions. Other environmental conditions in the leaf cuvette were as follows: temperature 20 +/- 0.5 °C; VPD 0.7 +/- 0.2 kPa; CO₂ of input air 360 +/-15 ppm.

4.1.2 Source/Platform

Branch samples were harvested in the early morning using a shotgun and transported to the laboratory in Thompson for gas exchange measurement.

4.1.3 Source/Platform Mission Objectives

The mission objectives were:

• To obtain the response curves of photosynthesis to ambient and intercellular CO₂ concentration, temperature, PAR, leaf-to-air VPD, and shoot water potential.

To examine interspecific differences in photosynthetic response to CO₂, temperature, PAR,

leaf-to-air VPD and shoot water potential.

• To examine seasonal variations in photosynthetic response to CO₂, temperature, PAR, leaf-to-air VPD, and shoot water potential.

4.1.4 Key Variables

Net photosynthesis, stomatal conductance, transpiration, ambient and intercellular CO₂ concentration, temperature, PAR flux density, VPD, water potential.

4.1.5 Principles of Operation

The stems of samples were connected to a water reservoir during the measurement to keep a continuous supply of water to the foliage. Independent samples were used for each temperature level, and each sample was measured for two CO₂ levels. Samples were exposed to saturated light for 2 hours prior to measurement to induce photosynthetic activity and stomatal opening.

Upper-canopy branch samples were harvested using a shotgun and were immediately recut under water. The samples were then transported to the laboratory for gas exchange measurement. The cut

surfaces of the branches were submerged in water during transportation (30 to 50 min).

In the laboratory, the branches were taken out of the water and the cut surfaces were dried and sealed using silicon grease. The branches were then exposed to light and let transpire freely. At certain time intervals, the gas exchange of the branches (two at a time) was measured. The water potential of the branches was measured immediately after the gas exchange measurement.

Gas exchange was measured at saturated light conditions. Other environmental conditions in the leaf cuvette were as follows: temperature 20 +/- 0.5 °C; VPD 0.7 +/- 0.2 kPa; CO₂ of input air 360

+/-15 ppm.

4.1.6 Sensor/Instrument Measurement Geometry

All samples were taken from the upper third of the forest canopy. Efforts were made to keep the amount of foliage relatively consistent from sample to sample. The leaf chamber for the measurement is about 1,300 in³.

4.1.7 Manufacturer of Sensor/Instrument

LI-6200 portable gas exchange system LI-COR P.O. Box 4425 4421 Superior St. Lincoln, NE 68504 (800) 447-3576

Leaf area measurement system/optical image analysis system (AgVision, monochrome system, root and leaf analysis)

Decagon Devices, Inc. P.O. Box 835 Pullman, WA 99163 (800) 755-2751

Pressure Chamber, Model 610 PMS Instrument Co. 480 SW Airport Avenue Corvallis, OR 97333 (503) 752-7926

4.2 Calibration

The LI-COR 6262 gas analyzer was calibrated using a standard gas at the beginning of each field campaign. The standard gas had been calibrated against the prime CO₂ standard in the NSA laboratory in Thompson, Manitoba, using gas chromatography technique. The stability of gas exchange and the reliability of the cut branch technique were also tested (see Dang et al., 1997a, for details).

4.2.1 Specifications

The weighing balance was accurate to within 0.0001 g. The leaf area system was accurate to within 1%. The gas exchange system was accurate to 1 ppm CO₂.

The shape factor used for black spruce was 4, in accordance with the BOREAS Experiment Plan, Appendix K, Version 3.0. Based on observations of two cross-sections of two needles per fascicle for five fascicles for six jack pine trees from Thompson, Manitoba, an average shape factor of 4.59 (+/-, 0.07) was calculated.

4.2.1.1 Tolerance

No tolerance level was set for these measurements.

4.2.2 Frequency of Calibration

The LI-COR 6262 IRGA was calibrated at the beginning of each IFC.

4.2.3 Other Calibration Information

Calibrations were performed according to each manufacturer's instructions.

5. Data Acquisition Methods

Upper-canopy branch samples were harvested using a shotgun and were immediately recut under water. The samples were then transported to the laboratory for gas exchange measurement. The cut surfaces of the branches were submerged in water during transport (30 to 50 min) and in the laboratory. Measurements for each species generally took 6 to 10 hours.

CO₂ concentration variation:

All samples were kept in the dark, but prior to measurement, samples were exposed to saturated light for 2 hours to induce stomatal opening and photosynthetic activity. Steady-state readings were taken at each CO₂ level, and an independent set of two samples was used for each two CO₂ levels. Ambient CO₂ varied from 50 to 900 ppm.

To keep a continued supply of water to the branch, the cut surface was kept in contact with water during the entire course of measurement. Saturated light was supplied using two 1,000-watt high-pressure sodium lamps. Other environmental conditions were as follows: temperature = 20 + 0.5 °C; VPD = 0.7 + 0.1 kPa; CO₂ = 360 + 0.2 ppm.

Temperature variation:

The air temperature inside the leaf cuvette was controlled using a radiator that was driven by a temperature-controlled water bath. The vapor pressure inside the cuvette was controlled by passing water vapor-saturated air through a condenser whose temperature was controlled using another water bath.

Saturated light for the measurement was supplied using two 1,000-watt high-pressure sodium lamps. VPD of the air was controlled at a relatively constant level except at temperatures below $10\,^{\circ}$ C, when there were some technical difficulties in getting a low enough vapor pressure to maintain the desired VPD. Input CO₂ concentration was controlled at 360 (+/- 15 ppm).

An independent set of two branches was used for each temperature. The branches were from four different trees and were mixed randomly.

PAR variations:

The measurements started from the highest PAR level and proceeded to darkness. Steady-state readings were taken at each light level. The light source was two 1,000-watt high-pressure sodium lamps. Different levels of light were achieved by using different neutral density filters. The environmental conditions inside the leaf cuvette were as follows: temperature 20 +/- 0.5 °C; VPD 0.7 +/- 0.2 kPa; CO₂ 360 +/- 15 ppm.

Vapor pressure variation:

All samples were kept in the dark, but prior to measurement, samples were exposed to saturated light for 2 hours to induce stomatal opening and photosynthetic activity. The stability and reliability of the cut-branch technique were tested. Stable measurements for at least 24 hours are possible. Steady-state readings were taken at each VPD level.

An independent set of two samples was used for each VPD level. To keep a continued supply of water to the branch, the cut surface was kept in contact with water during the entire course of

measurement.

Different VPD levels were achieved by regulating the water vapor pressure of the input air stream to the leaf cuvette. Saturated light was supplied using two 1,000-watt high-pressure sodium lamps. The CO_2 concentration in the input air was 360 (+/- 15 ppm). Measurements were taken at three temperatures (15, 25, and 35 °C) in IFC-1, two temperatures (25 and 35 °C) in IFC-2, and at 25 °C only in IFC-3.

Shoot water potential variations:

In the laboratory, the branches were taken out of the water and the cut surfaces were dried and sealed using silicon grease. The branches were then exposed to light and let transpire freely. At certain time intervals, the gas exchange of the branches (two at a time) was measured. The water potential of the branches was measured immediately after the gas exchange measurement.

Gas exchange was measured at saturated light conditions. Other environmental conditions in the leaf cuvette were as follows: temperature 20 +/- 0.5 °C; VPD 0.7 +/- 0.2 kPa; CO₂ of input air 360 +/-15 ppm.

6. Observations

6.1 Data Notes

Three to four leaves per sample for aspen.

6.2 Field Notes

Samples were taken from trees of relatively consistent vigor. See pages 2-23 and 2-24 in the BOREAS Experiment Plan, Version 3.0, for a description of site conditions.

7. Data Description

7.1 Spatial Characteristics

7.1.1 Spatial Coverage

At each site, branch samples were taken from four different trees that were at least 10 m apart from one another. Sampling was done within a 100 m² area. The North American Datum of 1983 (NAD83) coordinates for each site are:

NSA-OJP flux tower site, Lat/Long:55.92842°N, 98.62396°W; UTM Zone 14, N:6198176.3, E:523496.2

• NSA-OASP canopy access tower site (auxiliary site number T2Q6A, BOREAS Experiment Plan, Version 3), Lat/Long 55.88691°N, 98.67479°W; UTM Zone 14, N: 6193540.7, E: 520342

NSA-OBS flux tower site, Lat/Long: 55.88007°N, 98.48139°W; UTM Zone 14, N: 6192853.4 E: 532444.5

7.1.2 Spatial Coverage Map

None.

7.1.3 Spatial Resolution

These data are point source measurements from the sampled trees.

7.1.4 Projection

Not applicable.

7.1.5 Grid Description

Not applicable.

7.2 Temporal Characteristics

7.2.1 Temporal Coverage

All data were collected between 24-May-1994 and 19-Sep-1994. Samples were taken between 6:00 and 7:00 a.m. Measurements in the laboratory generally took 6 to 8 hours. An independent data set was taken during each of the three field campaigns. The specific dates for each data set are given in the data table.

7.2.2 Temporal Coverage Map

CO2 concentration:

Site	Sample	Dates (da	ay-month)	1994
NSA-OBS	04-JUN,	09-AUG,	06-SEP	
NSA-OJP	02-JUN,	06-AUG,	07-SEP	
NSA-OA	01-JUN.	07-AUG.	30-AUG	

Temperature:						
Site	Sample I	Dates (da	ay-month)	1994		
NSA-OBS	23-MAY,	26-JUL,	15-SEP			
NSA-OJP	13-MAY,	27-JUL,	16-SEP			
NSA-OA	14-JUN,	28-JUL,	10-SEP			

PAR:

Site	Sample I	Dates (da	ay-month)	1994
NSA-OBS	25-MAY,	23-JUL,	14-SEP	
NSA-OJP	26-MAY,	24-JUL,	13-SEP	
NSA-OA	10-JUN,	25-JUL,	09-SEP	

VPD:

Site	Sample I	Dates (d	ay-month;) 1994		
NSA-OBS	30-MAY,	25-MAY,	28-MAY,	29-AUG,	30-AUG,	12-SEP
NSA-OJP	24-MAY,	27-MAY,	29-MAY,	01-AUG,	03-AUG,	17-SEP

Water Shoot Potential:

```
Site Sample Dates (day-month) 1994
NSA-OBS 23-JUL, 06-SEP
NSA-OJP 02-AUG, 07-SEP
```

7.2.3 Temporal Resolution

The measurements can be considered to be single point in time measurements since the same trees were not repeatedly sampled.

7.3 Data Characteristics

7.3.1 Parameter/Variable

The parameters contained in the data files on the CD-ROM are:

SITE NAME SUB SITE START_DATE END DATE SPECIES PARAM VARIED LEAF TEMP AIR_TEMP CO2 CONC TRANSPIRATION RATE PHOTOSYNTHETIC RATE DOWN PPFD INTERCELL CO2 CONC STOMATAL_CONDUCT_CO2 WATER_USE_EFF WATER POTENTIAL VAPOR_PRESS_DEFICIT VAPOR PRESS DEFICIT AIR TEMP CRTFCN CODE REVISION_DATE

7.3.2 Variable Description/Definition

The descriptions of the parameters contained in the data files on the CD-ROM are:

Column Name	Description
SITE_NAME	The identifier assigned to the site by BOREAS, in the format SSS-TTT-CCCCC, where SSS identifies the portion of the study area: NSA, SSA, REG, TRN, and TTT identifies the cover type for the site, 999 if unknown, and CCCCC is the identifier for site, exactly what it means will vary with site type.
SUB_SITE	The identifier assigned to the sub-site by BOREAS, in the format GGGGG-IIIII, where GGGGG is the group associated with the sub-site instrument, e.g. HYD06 or STAFF, and IIIII is the identifier for sub-site, often this will refer to an instrument.
START_DATE	The date on which the collection of the referenced data commenced.
END_DATE	The date on which the collection of the referenced data was terminated.
SPECIES	Botanical (Latin) name of the species (Genus species).
PARAM_VARIED	The parameter varied to study photosynthetic response to BOREAL tree species. i.e. CO2

CONCENTRATION=CO2 varied; LIGHT=light varied; TEMPERATURE=temperature varied; VAPOR PRESS DEFICIT = vapor pressure deficit varied; WATER POTENTIAL = Water potential varied; DARK RESPIRATION=temperature varied in dark; HUMIDITY=humidity varied; CONSTANT CONDITIONS=nothing varied; STOMATAL MODEL=many things varied for model calibration. LEAF TEMP The measured leaf or shoot temperature AIR TEMP The measured air temperature. CO2 CONC CO2 concentration. TRANSPIRATION RATE Transpiration rate (E) Measured Net Photosynthesis
The downward photosynthetic photon flux density. PHOTOSYNTHETIC RATE DOWN PPFD INTERCELL CO2 CONC Intercellular CO2 concentration STOMATAL CONDUCT CO2 Stomatal conductance to CO2 (gs) WATER USE EFF Water use efficiency WATER POTENTIAL Water Potential VAPOR_PRESS_DEFICIT Vapor Pressure Deficit (VPD) VAPOR_PRESS_DEFICIT_AIR_TEMP The air temperature when vapor pressure deficit was varied. CRTFCN CODE The BOREAS certification level of the data. Examples are CPI (Checked by PI), CGR (Certified by Group), PRE (Preliminary), and CPI-??? (CPI but questionable). REVISION DATE The most recent date when the information in the referenced data base table record was revised.

7.3.3 Unit of Measurement

The measurement units for the parameters contained in the data files on the CD-ROM are:

Column Name	Units
SITE NAME	[none]
SUB SITE	[none]
START DATE	[DD-MON-YY]
END DATE	[DD-MON-YY]
SPECIES	[none]
PARAM_VARIED	[none]
LEAF_TEMP	[degrees Celsius]
AIR_TEMP	[degrees Celsius]
CO2_CONC	[parts per million]
TRANSPIRATION_RATE	<pre>[millimoles] [meter^-2] [second^-1]</pre>
PHOTOSYNTHETIC_RATE	<pre>[micromoles] [meter^-2] [second^-1]</pre>
DOWN_PPFD	<pre>[micromoles] [meter^-2] [second^-1]</pre>
INTERCELL_CO2_CONC	[parts per million]
STOMATAL_CONDUCT_CO2	<pre>[millimoles] [meter^-2] [second^-1]</pre>
WATER_USE_EFF	[micromoles CO2][millimole H2O^-1]
WATER_POTENTIAL	[megaPascals]
VAPOR_PRESS_DEFICIT	[kiloPascals]
VAPOR_PRESS_DEFICIT_AIR_TEMP	[degrees Celsius]
CRTFCN_CODE	[none]
REVISION_DATE	[DD-MON-YY]

7.3.4 Data Source

The sources of the parameter values contained in the data files on the CD-ROM are:

Column Name	Data Source
SITE_NAME SUB_SITE START_DATE END_DATE	BORIS Designation BORIS Designation BORIS Designation BORIS Designation
SPECIES	Human Observer Human Observer
PARAM_VARIED LEAF_TEMP AIR TEMP	Thermometer Thermometer
CO2_CONC	Laboratory Equipment
TRANSPIRATION_RATE	Laboratory Equipment
PHOTOSYNTHETIC_RATE	Laboratory Equipment
DOWN_PPFD INTERCELL CO2 CONC	PPFD Sensor Laboratory Equipment
STOMATAL_CONDUCT_CO2	Laboratory Equipment
WATER_USE_EFF	Laboratory Equipment
WATER_POTENTIAL	Laboratory Equipment
VAPOR_PRESS_DEFICIT	Laboratory Equipment
VAPOR_PRESS_DEFICIT_AIR_TEMP CRTFCN CODE	Thermometer BORIS Designation
REVISION_DATE	BORIS Designation

7.3.5 Data Range

The following table gives information about the parameter values found in the data files on the CD-ROM.

Column Name	Minimum Data Value	Maximum Data Value		Data		Data Not Cllctd
SITE NAME	NSA-90A-9TETR	NSA-OJP-FLXTR	None	None	None	None
	9TE09-PHR01	9TE09-PHR01	None	None	None	None
	24-MAY-94	30-AUG-94	None	None	None	None
	16-JUN-94	19-SEP-94	None	None	None	None
SPECIES	N/A	N/A	None	None	None	None
PARAM VARIED .	N/A	N/A	None	None	None	None
LEAF TEMP	-4.94	44.22	None	None	None	None
AIR TEMP	-6.94	43.9	None	None	None	None
CO2 CONC	47.53	872	None	None	None	None
TRANSPIRATION_RATE	.004843	8	None	None	None	None
PHOTOSYNTHETIC_RATE	-4.55426	25.8703	None	None	None	None
DOWN_PPFD	0	1452	None	None	None	None
INTERCELL_CO2_CONC	31.526	667.147	None	None	None	None
STOMATAL_CONDUCT_CO2	.5891	149.8545	None	None	None	None
WATER_USE_EFF	-20.842	85.604	None	None	None	None
WATER POTENTIAL	.15	3.95	None	None	None	Blank
VAPOR_PRESS_DEFICIT	. 04	5.368694	None	None	None	None
VAPOR_PRESS_DEFICIT_		25	None	None	None	Blank
AIR_TEMP CRTFCN_CODE	CPT	CPI	None	None	None	None
REVISION_DATE	03-SEP-96			None	None	None

Minimum Data Value -- The minimum value found in the column. Maximum Data Value -- The maximum value found in the column. Missng Data Value -- The value that indicates missing data. This is used to indicate that an attempt was made to determine the parameter value, but the attempt was unsuccessful. -- The value that indicates unreliable data. This is used Unrel Data Value to indicate an attempt was made to determine the parameter value, but the value was deemed to be unreliable by the analysis personnel. Below Detect Limit -- The value that indicates parameter values below the instruments detection limits. This is used to indicate that an attempt was made to determine the parameter value, but the analysis personnel determined that the parameter value was below the detection limit of the instrumentation. Data Not Cllctd -- This value indicates that no attempt was made to determine the parameter value. This usually indicates that BORIS combined several similar but not identical data sets into the same data base table but this particular science team did not measure that parameter.

Blank -- Indicates that blank spaces are used to denote that type of value.

N/A -- Indicates that the value is not applicable to the respective column.

None -- Indicates that no values of that sort were found in the column.

7.4 Sample Data Record

The following is a sample of the first few records from the data table on the CD-ROM:

SITE_NAME, SUB_SITE, START_DATE, END_DATE, SPECIES, PARAM_VARIED, LEAF_TEMP, AIR_TEMP, CO2_CONC, TRANSPIRATION_RATE, PHOTOSYNTHETIC_RATE, DOWN_PPFD, INTERCELL_CO2_CONC, STOMATAL_CONDUCT_CO2, WATER_USE_EFF, WATER_POTENTIAL, VAPOR_PRESS_DEFICIT, VAPOR_PRESS_DEFICIT_AIR_TEMP, CRTFCN_CODE, REVISION_DATE
'NSA-OBS-FLXTR', '9TE09-PHR01', 24-MAY-94, 16-JUN-94, 'Picea mariana', 'LIGHT', 19.8, 19.8, 359.3, .081231, -.58655, 0.0, 424.4152, 7.836619, -7.22077, .618614, , 'CPI', 03-SEP-96
'NSA-OBS-FLXTR', '9TE09-PHR01', 24-MAY-94, 16-JUN-94, 'Picea mariana', 'LIGHT', 20.3, 20.3, 333.1, .195906, 1.968549, 1007.0, 254.9745, 20.48158, 10.04844, .570307, , 'CPI', 03-SEP-96

8. Data Organization

8.1 Data Granularity

The smallest unit of orderable data is data collected on one day at one site.

8.2 Data Format(s)

The Compact Disk-Read-Only Memory (CD-ROM) files contain American Standard Code for Information Interchange (ASCII) numerical and character fields of varying length separated by commas. The character fields are enclosed with single apostrophe marks. There are no spaces between the fields.

Each data file on the CD-ROM has four header lines of Hyper-Text Markup Language (HTML) code at the top. When viewed with a Web browser, this code displays header information (data set title, location, date, acknowledgments, etc.) and a series of HTML links to associated data files and related data sets. Line 5 of each data file is a list of the column names, and line 6 and following lines contain the actual data.

9. Data Manipulations

9.1 Formulae

9.1.1 Derivation Techniques and Algorithms

A, E, gs-CO₂, and Ci were calculated according to von Caemmerer and Farquhar (1981), Planta 153: 376-387.

WUE = A/E

where: WUE = photosynthetic water use efficiency

A = net photosynthesis E = transpiration rate

VPD = VPsat - Vpamt

where: VPD = vapor pressure difference

VPsat is saturated vapor pressure in the chamber VPamt is measured vapor pressure in the chamber

9.2 Data Processing Sequence

9.2.1 Processing Steps

Data were recorded automatically by a computer and also printed on a printer. Subsequent calculations of different parameters were performed using MS Excel for Windows 5.0.

BOREAS Information System (BORIS) staff processed the data by:

- Reviewing the initial data files and loading them online for BOREAS team access.
- Designing relational data base tables to inventory and store the data.
- Loading the data into the relational data base tables.
- Performing the following conversions on measurements into System International (SI) units: -Changing PAR flux from (mol/m²/s) to DOWN_PPFD (μmol/m²/s)
- Working with the Terrestrial Ecology (TE)-09 team to document the data set.
- Extracting the standardized data into logical files.

9.2.2 Processing Changes

None.

9.3 Calculations

A, E, gs-CO₂, and Ci were calculated according to von Caemmerer and Farquhar (1981), Planta 153: 376-387.

WUE = A/EVPD = VPsat - Vpamt

where: VPsat and VPamt are saturated vapor pressure and measured vapor pressure in the chamber.

9.3.1 Special Corrections/Adjustments

None.

9.3.2 Calculated Variables

A, E, gs-CO₂, and Ci were calculated according to von Caemmerer and Farquhar (1981), Planta 153: 376-387.

WUE = A/E VPD = VPsat - Vpamt

where: VPsat and VPamt are saturated vapor pressure and measured vapor pressure in the chamber.

9.4 Graphs and Plots

Net photosynthesis versus ambient and internal CO₂ concentration. A, gs, WUE vs. P

10. Errors

10.1 Sources of Error

During photosynthetic response to temperature differences, condensation sometimes formed on the radiator inside the cuvette when the temperature went below 10 °C.

Possible genetic differences between trees and possible differences in physiological conditions between branches could cause inconsistencies in the data.

There are no other known sources of error.

10.2 Quality Assessment

Please contact Dr. Hank Margolis and Dr. Qinglai Dang if these data are used for publication (see Section 2.3, Contact Information).

10.2.1 Data Validation by Source

After each measurement, the sample was removed from the leaf cuvette and a base measurement (i.e., when cuvette contained no sample) was taken. The previous measurement was adjusted by this base value, if necessary.

10.2.2 Confidence Level/Accuracy Judgment

No statistical confidence level is yet available. However, the investigators are very confident that these data are reliable. Results are consistent with field measurements.

10.2.3 Measurement Error for Parameters

Unknown.

10.2.4 Additional Quality Assessments

Calculated results were plotted, and the patterns were examined. Obvious outliers (determined visually) were eliminated from the data set.

10.2.5 Data Verification by Data Center

Data were examined for general consistency and clarity

11. Notes

11.1 Limitations of the Data

None given.

11.2 Known Problems with the Data

None.

11.3 Usage Guidance

Parameters derived from this data set will be more applicable to aggregated foliage on the shoot as a whole than to individual needles or leaves.

11.4 Other Relevant Information

None.

12. Application of the Data Set

Data can be used to examine the influence of different factors on the photosynthetic process.

13. Future Modifications and Plans

None.

14. Software

14.1 Software Description

Calculations were performed using MS Excel for Windows 5.0.

14.2 Software Access

Contact Microsoft Corp.

15. Data Access

The NSA photosynthetic response data are available from the Earth Observing System Data and Information System (EOSDIS) Oak Ridge National Laboratory (ORNL) Distributed Active Archive Center (DAAC).

15.1 Contact Information

For BOREAS data and documentation please contact:

ORNL DAAC User Services
Oak Ridge National Laboratory
P.O. Box 2008 MS-6407
Oak Ridge, TN 37831-6407

Phone: (423) 241-3952 Fax: (423) 574-4665

E-mail: ornldaac@ornl.gov or ornl@eos.nasa.gov

15.2 Data Center Identification

Earth Observing System Data and Information System (EOSDIS) Oak Ridge National Laboratory (ORNL) Distributed Active Archive Center (DAAC) for Biogeochemical Dynamics http://www-eosdis.ornl.gov/.

15.3 Procedures for Obtaining Data

Users may obtain data directly through the ORNL DAAC online search and order system [http://www-eosdis.ornl.gov/] and the anonymous FTP site [ftp://www-eosdis.ornl.gov/data/] or by contacting User Services by electronic mail, telephone, fax, letter, or personal visit using the contact information in Section 15.1.

15.4 Data Center Status/Plans

The ORNL DAAC is the primary source for BOREAS field measurement, image, GIS, and hardcopy data products. The BOREAS CD-ROM and data referenced or listed in inventories on the CD-ROM are available from the ORNL DAAC.

16. Output Products and Availability

16.1 Tape Products

None.

16.2 Film Products

None.

16.3 Other Products

These data are available on the BOREAS CD-ROM series.

17. References

17.1 Platform/Sensor/Instrument/Data Processing Documentation

LI-COR 6262 Infrared gas analyzer manual.

17.2 Journal Articles and Study Reports

Dang, Q.L., H. Margolis, M.R. Coyea, M. Sy, and G.J. Collatz. 1997a. Regulation of branch-level gas exchange of boreal trees: roles of shoot water potential and vapor pressure difference. Tree Physiology, BOREAS Special Issue 17(8/9):521-535.

Dang, Q.L., H. Margolis, G.J. Collatz, et al. Parameterization and testing of a coupled photosynthesis stomatal conductance model for the boreal forest. Tree Physiology (in preparation).

Newcomer, J., D. Landis, S. Conrad, S. Curd, K. Huemmrich, D. Knapp, A. Morrell, J. Nickeson, A. Papagno, D. Rinker, R. Strub, T. Twine, F. Hall, and P. Sellers, eds. 2000. Collected Data of The Boreal Ecosystem-Atmosphere Study. NASA. CD-ROM.

Sellers, P. and F. Hall. 1994. Boreal Ecosystem-Atmosphere Study: Experiment Plan. Version 1994-3.0, NASA BOREAS Report (EXPLAN 94).

Sellers, P. and F. Hall. 1996. Boreal Ecosystem-Atmosphere Study: Experiment Plan. Version 1996-2.0, NASA BOREAS Report (EXPLAN 96).

Sellers, P., F. Hall, and K.F. Huemmrich. 1996. Boreal Ecosystem-Atmosphere Study: 1994 Operations. NASA BOREAS Report (OPS DOC 94).

Sellers, P., F. Hall, and K.F. Huemmrich. 1997. Boreal Ecosystem-Atmosphere Study: 1996 Operations. NASA BOREAS Report (OPS DOC 96).

Sellers, P., F. Hall, H. Margolis, B. Kelly, D. Baldocchi, G. den Hartog, J. Cihlar, M.G. Ryan, B. Goodison, P. Crill, K.J. Ranson, D. Lettenmaier, and D.E. Wickland. 1995. The boreal ecosystem-atmosphere study (BOREAS): an overview and early results from the 1994 field year. Bulletin of the American Meteorological Society. 76(9):1549-1577.

Sellers, P.J., F.G. Hall, R.D. Kelly, A. Black, D. Baldocchi, J. Berry, M. Ryan, K.J. Ranson, P.M. Crill, D.P. Lettenmaier, H. Margolis, J. Cihlar, J. Newcomer, D. Fitzjarrald, P.G. Jarvis, S.T. Gower, D. Halliwell, D. Williams, B. Goodison, D.E. Wickland, and F.E. Guertin. 1997. BOREAS in 1997: Experiment Overview, Scientific Results and Future Directions. Journal of Geophysical Research 102(D24): 28,731-28,770.

von Caemmerer, S. and G.D. Farquhar. 1981. Some relationships between biochemistry of photosynthesis and the gas exchange of leaves. Planta 153: 376-387.

17.3 Archive/DBMS Usage Documentation None.

18. Glossary of Terms

```
- net photosynthesis (\mumol CO<sub>2</sub>/m<sup>2</sup>/s)
A-Ci curve - photosynthetic response to CO2
             - intercellular CO2 concentration (ppm)
Ci
            - ambient CO2 concentration (ppm)
CO<sub>2</sub>
             - transpiration rate (mmol H<sub>2</sub>O/m<sup>2</sup>/s)
qs CO<sub>2</sub>
           - stomatal conductance to CO<sub>2</sub> (mmol/m<sup>2</sup>/s)
            - shoot water potential (MPa)
Tleaf
          leaf temperature (°C)
         - air temperature (°C)
Tair
            - photosynthetic water use efficiency (mmol CO_2/mol\ H_2O)
WUE
```

19. List of Acronyms

```
ASCII - American Standard Code for Information Interchange
BOREAS - BOReal Ecosystem-Atmosphere Study
BORIS - BOREAS Information System
CD-ROM - Compact Disk-Read-Only Memory
CGI - Certified by Group
CPI - Checked by Principal Investigator
DAAC - Distributed Active Archive Center
EOS - Earth Observing System
EOSDIS - EOS Data and Information System
GIS - Geographic Information System
GSFC - Goddard Space Flight Center
HTML - Hyper-Text Markup Language
```

IFC - Intensive Field Campaign - Infrared Gas Analyzer IRGA NAD83 - North American Datum of 1983 NASA - National Aeronautics and Space Administration NSA - Northern Study Area ORNL - Oak Ridge National Laboratory - Prince Albert National Park PANP - Photosynthetically Active Radiation PAR PRE - Preliminary SSA - Southern Study Area URL - Uniform Resource Locator - Universal Transverse Mercator UTM VPD - Vapor Pressure Difference (kPa)

20. Document Information

20.1 Document Revision Date

Written: 12-Mar-1996 Last updated: 20-Apr-1999

20.2 Document Review Dates

BORIS Review: 22-Apr-1997 Science Review: 05-Nov-1997

20.3 Document ID

20.4 Citation

When using these data, please contact one of the individuals listed in Section 2.3 as well as citations of relevant papers in Section 17.2.

If using data from the BOREAS CD-ROM series, also reference the data as:

H. Margolis, "Relationship Between Measures of Absorbed and Reflected Radiation and the Photosynthetic Capacity of Boreal Forest Canopies and Understories." In Collected Data of The Boreal Ecosystem-Atmosphere Study. Eds. J. Newcomer, D. Landis, S. Conrad, S. Curd, K. Huemmrich, D. Knapp, A. Morrell, J. Nickeson, A. Papagno, D. Rinker, R. Strub, T. Twine, F. Hall, and P. Sellers. CD-ROM. NASA, 2000.

Also, cite the BOREAS CD-ROM set as:

Newcomer, J., D. Landis, S. Conrad, S. Curd, K. Huemmrich, D. Knapp, A. Morrell, J. Nickeson, A. Papagno, D. Rinker, R. Strub, T. Twine, F. Hall, and P. Sellers, eds. Collected Data of The Boreal Ecosystem-Atmosphere Study. NASA. CD-ROM. NASA, 2000.

20.5 Document Curator

20.6 Document URL

	•	

Form Approved REPORT DOCUMENTATION PAGE

			OMB No. 0704-0188
gathering and maintaining the data needed, and co	mpleting and reviewing the collection of reducing this burden, to Washington Hea	information. Send comments reg douarters Services, Directorate f	eviewing instructions, searching existing data sources, parding this burden estimate or any other aspect of this or Information Operations and Reports, 1215 Jefferson Project (0704-0188), Washington, DC 20503.
1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE October 2000	3. REPORT TYPE AN Technical Me	
4. TITLE AND SUBTITLE			5. FUNDING NUMBERS
Technical Report Series on the R	oreal Ecosystem_Atmosphe	ore Study (RODEAS)	

BOREAS TE-9 NSA Photosynthetic Response Data 923

RTOP: 923-462-33-01 6. AUTHOR(S) Qinglai Dang, Hank Margolis, and Marie Coyea

Forrest G. Hall and Shelaine Curd, Editors

7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS (ES) 8. PEFORMING ORGANIZATION REPORT NUMBER

Goddard Space Flight Center 2000-03136-0 Greenbelt, Maryland 20771

9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS (ES) 10. SPONSORING / MONITORING AGENCY REPORT NUMBER

National Aeronautics and Space Administration TM-2000-209891 Washington, DC 20546-0001 Vol. 157

11. SUPPLEMENTARY NOTES Q. Dang: Lakehead University, Thunder Bay, Canada; H. Margolis and M. Coyea: Université Laval, Sainte-Foy, Quebec, Canada; S. Curd: Raytheon ITSS, NASA Goddard Space Flight Center,

Greenbelt, Maryland 12a. DISTRIBUTION / AVAILABILITY STATEMENT 12b. DISTRIBUTION CODE

Unclassified—Unlimited Subject Category: 43 Report available from the NASA Center for AeroSpace Information, 7121 Standard Drive, Hanover, MD 21076-1320. (301) 621-0390.

13. ABSTRACT (Maximum 200 words)

The BOREAS TE-9 team collected several data sets related to chemical and photosynthetic properties of leaves. This data set describes (1) the response of leaf and shoot-level photosynthesis to ambient and intercellular CO, concentration, temperature, and incident PAR for black spruce, jack pine, and aspen during the three IFCs in 1994 in the NSA; (2) the response of stomatal conductance to vapor pressure difference throughout the growing season of 1994; and (3) a range of shoot water potentials (controlled in the laboratory) for black spruce and jack pine. The data are available in tabular ASCII files.

14. SUBJECT TERMS BOREAS, terrestrial eco	15. NUMBER OF PAGES 18 16. PRICE CODE		
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT UL

	•	